

BJPpsych

The British Journal of Psychiatry

Involuntary out-patient commitment and reduction of violent behaviour in persons with severe mental illness

JEFFREY W. SWANSON, MARVIN S. SWARTZ, H. RYAN WAGNER, BARBARA J. BURNS, RANDY BORUM and VIRGINIA A. HIDAY

BJP 2000, 176:324-331.

Access the most recent version at DOI: [10.1192/bjp.176.4.324](https://doi.org/10.1192/bjp.176.4.324)

References

This article cites 0 articles, 0 of which you can access for free at:
<http://bjp.rcpsych.org/content/176/4/324#BIBL>

Reprints/ permissions

To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk

You can respond to this article at

<http://bjp.rcpsych.org/cgi/eletter-submit/176/4/324>

Downloaded from

<http://bjp.rcpsych.org/> on March 7, 2012
Published by The Royal College of Psychiatrists

Involuntary out-patient commitment and reduction of violent behaviour in persons with severe mental illness[†]

JEFFREY W. SWANSON, MARVIN S. SWARTZ, RANDY BORUM, VIRGINIA A. HIDAY, H. RYAN WAGNER and BARBARA J. BURNS

Background Violent behaviour among persons with severe mental illness (SMI) causes public concern and is associated with illness relapse, hospital recidivism and poor outcomes in community-based treatment.

Aims To test whether involuntary out-patient commitment (OPC) may help to reduce the incidence of violence among persons with SMI.

Method One-year randomised trial of the effectiveness of OPC in 262 subjects with psychotic or major mood disorders and a history of hospital recidivism. Involuntarily hospitalised subjects awaiting OPC were randomly assigned to release or court-ordered treatment after discharge. Those with a recent history of serious assault remained under OPC until expiry of the court order (up to 90 days); then OPC orders were renewed at clinical/court discretion. Control subjects had no OPC. Four-monthly follow-up interviews with subject, case manager and collateral informant took place and service records were collected.

Results A significantly lower incidence of violent behaviour occurred in subjects with ≥ 6 months' OPC. Lowest risk of violence was associated with extended OPC combined with regular out-patient services, adherence to prescribed medications and no substance misuse.

Conclusions OPC may significantly reduce risk of violent behaviour in persons with SMI, in part by improving adherence to medications while diminishing substance misuse.

Declaration of interest Funded by the National Institute of Mental Health.

Violent behaviour in persons with psychiatric disorder causes great public concern and is recognised by mental health clinicians as a significant problem associated with illness relapse, hospital recidivism and poor outcomes in community-based treatment (Monahan & Steadman, 1994; Mullen, 1997). Involuntary out-patient commitment (OPC) is a legal intervention designed to benefit individuals with severe mental illness (SMI) who need ongoing psychiatric care and support to prevent dangerous relapse, but who are reluctant or unable to follow through with community-based treatment. Proponents of OPC assert that it may improve treatment adherence and may also act as a lever on a mental health service system: mobilising supportive services and motivating clinical vigilance. This paper addresses the question of whether OPC can help to prevent violent behaviour in psychiatric populations with a history of 'revolving-door' hospital admissions often associated with dangerousness. Findings are presented from the first randomised study of the effectiveness of OPC combined with case management (Swartz *et al*, 1999).

Background

Serious violent acts committed by individuals with mental illnesses are statistically rare events (Monahan, 1992). Still, the potential for violence in a proportion of persons with severe, persistent and often untreated psychiatric disorders stimulates public fear, prevents general acceptance and inclusion of persons with psychiatric disabilities and limits normalisation and effectiveness of community-based mental health services. The risk of violence creates dilemmas in the clinical realm by interrupting community tenure and continuity of care, in the legal realm by increasing concerns about professional liability, and in the public realm by heightening fear and stigma associated with mental

illness (Link *et al*, 1987; Angermeyer & Matschinger, 1996; Borum *et al*, 1996; Simon, 1998).

With the advent of managed care in both public and private mental health systems, and with clinicians increasingly held liable for the behaviour of patients inadequately treated, concerns about the risk of violence have increased (Cuffel, 1997; Simon, 1998). Individuals suffering from SMI with a history of violence are disproportionately high users of the most expensive mental health services in the most restrictive settings (e.g. involuntary in-patient treatment) (Borum *et al*, 1996). However, clinicians' efforts to prevent violence through conventional out-patient treatment are impeded: patients at risk often do not adhere to medication regimens or keep scheduled appointments, they may misuse substances and they tend to live in impoverished, dangerous environments with inadequate social support (Swartz *et al*, 1998b; Silver *et al*, 1999). Some studies have reported increased violent behaviour among mentally ill persons suffering from psychotic symptoms that involve perceived threat and override of internal cognitive controls (threat/control-override: TCO) (Link *et al*, 1998), and particularly among those with TCO symptoms and substance misuse who have not had recent contact with a community mental health provider (Swanson *et al*, 1997a).

Involuntary OPC is a promising legal intervention that may significantly reduce violent behaviour associated with these particular problems by improving compliance with medications that mitigate high-risk psychotic symptoms; by improving access to substance misuse treatment for persons with dual diagnoses; by increasing clinical surveillance; and by augmenting case management intensity, thus leveraging scarce resources in community care systems (Swanson *et al*, 1997b).

Out-patient commitment statutes exist in 35 states and the District of Columbia (Torrey & Kaplan, 1995). In North Carolina, where the present study was conducted, OPC statutes require compliance with recommended treatment, excluding forced medication in the out-patient setting. Under OPC, the responsible clinician may request that law officers escort the non-adherent patient to a community mental health centre for examination and hopeful persuasion to accept treatment. Evidence from naturalistic studies

[†]See editorial pp. 307-311, this issue.

(Fernandez & Nygard, 1990; Geller *et al*, 1998), as well as the current randomised study (Swartz *et al*, 1999), suggests that OPC may decrease hospital readmission rates and total days hospitalised. However, OPC is strongly opposed by some mental health consumers and mental health law advocates, who argue that coerced out-patient treatment infringes on civil liberties, extends unwarranted social control into the community and may actually drive people away from needed treatment (Mulvey *et al*, 1987; Stefan, 1987).

The potential of OPC to prevent violent behaviour warrants empirical investigation. The present study addresses the question of whether and how OPC reduces the risk of violence among people with SMI, using prospective longitudinal data from a sample of 262 people with psychotic or major mood disorders who were placed on OPC combined with case management in the Piedmont region of North Carolina.

METHOD

Study design and sample

Selection criteria

Subjects were screened sequentially from a population of involuntarily hospitalised patients who had been ordered to undergo a period of OPC upon discharge. Eligibility criteria for the study were: age 18 years or older; diagnosis of schizophrenia, schizoaffective disorder, other psychotic disorder or major affective disorder; duration of disorder of one year or more; significant functional impairment in activities of daily living; intensive treatment within the past two years; resident of one of nine counties participating in the study; and awaiting a period of court-ordered OPC. Legal criteria for OPC in North Carolina include SMI diagnosis and mental status limiting a person's ability to seek or comply voluntarily with treatment, and the likelihood that without treatment the person would predictably decompensate to a point of dangerousness or grave disability.

Study group assignment

By special arrangement with the court, subjects randomly assigned to a control group were released from OPC. Subjects in the experimental group, by law, received an initial period of OPC no longer than 90 days. Thereafter, the commitment order could be renewed for up to 180 days if a

psychiatrist and the court determined that the subject continued to meet the legal criteria for OPC. However, subjects in the control group received immunity from any OPC during the year of the study. All subjects received case management and other out-patient treatment at one of four participating area mental health programmes representing nine contiguous urban and rural counties. An exception to the randomisation procedure was necessary in the case of subjects with a history of serious assault involving weapon use or physical injury to another person within the preceding year. These subjects (the seriously violent group) were required to undergo at least the initial period of OPC as ordered. Renewals were left to the discretion of the clinician and court.

Refusal, attrition, and differences in length of time on OPC

Of the identified eligible patients, 12% refused consent to participate. Rates of refusal did not vary significantly by gender, race or diagnosis. Subjects over age 45 years were more likely to refuse than those under 45 years (14% *v.* 7%). The baseline sample consisted of 331 subjects. At the 12-month follow-up, 69 subjects (20.9%) had withdrawn or were lost to follow-up, with 262 remaining: 114 controls, 102 in the OPC group and 46 in the OPC violent group. Attrition did not differ significantly by group. There was no evidence of sample bias in renewal of OPC orders, except that subjects with a baseline history of medication non-compliance were somewhat more likely to receive extended OPC (renewed court orders); 40.0% *v.* 18.75%. (The potential implications of this difference for interpretation of the results will be discussed below.) Approximately one-third of subjects in both the OPC and violent OPC groups received overall more than 180 days of court-ordered treatment.

Data collection

At baseline, structured interviews were conducted with each subject and with a family member or other informant who knew the respondent well. Hospital records were reviewed for additional information regarding clinical history. Follow-up interviews were conducted every four months with the subjects, case manager and collateral informant. Out-patient service records and hospital admissions were recorded as well.

Measurement

Violence

Incidence of violence was assessed from three data sources. Subjects were asked every four months whether they had been picked up by police or arrested for physical assault on another person, had been in fights involving physical contact or had threatened someone with a weapon. Family members and case managers were asked comparable questions about the subject's behaviour. A composite index was constructed measuring whether at least one violent act was reported by any source during the year of the study (Swanson *et al*, 1999).

Psychiatric symptoms

Psychiatric symptoms were assessed using the Brief Symptom Inventory (BSI; Derogatis & Melisaratos, 1983). The BSI total score and a sub-scale for paranoid symptoms were used in the present analysis. An additional sub-scale was constructed from BSI items measuring TCO symptoms (Link *et al*, 1998).

Functional impairment

Functional impairment was assessed using the Global Assessment of Functioning scale (GAF), which is a measure of functional status and severity of psychiatric disturbance rated on a continuum of 0–100 from most to least impaired (Endicott *et al*, 1976).

Insight

Insight was measured by the Insight and Treatment Attitudes Questionnaire (ITAQ; McEvoy *et al*, 1989). The ITAQ assesses the ability of subjects with SMI to recognise their need for treatment.

Social support

Social support was assessed using a sub-scale of the Duke Social Support Scale (George *et al*, 1989), measuring respondents' subjective perception of their status and value in a social network, whether the network would provide help if needed and satisfaction with the quantity and quality of received support.

Substance use and misuse

Substance use and misuse were assessed by combining interview data from three sources and (at baseline) the hospital

record. Substance use was defined as drinking alcohol or using illicit drugs once to several times per month or more frequently. Substance misuse was defined as the occurrence of any problems related

to alcohol or drug use: problems with family, friends, job, police, physical health or any recorded diagnosis of psychoactive substance use disorder (Swartz *et al*, 1998a).

Table 1 One-year incidence of violent behaviour by sample characteristics

Baseline risk factors		<i>n</i>	% Violent
Age	18–39 years	138	48.55***
	40–65 years	124	26.61
Gender	Male	140	40.71
	Female	122	35.25
Race	African American	172	42.44†
	White/other	90	30.00
Marital status	Single	208	41.35*
	Married	54	25.93
Duke Social Support Scale (perceived social support)	Median or below	144	47.22***
	Above median	118	27.12
Education	High school graduate	165	38.79
	No High School	97	37.11
Annual income	\$6000 or less	137	40.15
	Above \$6000	125	36.00
Community of residence	Rural	108	30.56*
	Urban	154	43.51
Homeless at some time in preceding four months	No	208	33.17***
	Yes	54	57.41
Victim of crime during preceding four months	No	194	35.05†
	Yes	68	47.06
Primary discharge diagnosis	Schizophrenia	100	35.00
	Schizoaffective	58	41.38
	Other psychotic	21	33.33
	Bipolar disorder	72	41.67
	Major depression	11	36.36
Personality disorder (Axis 2 diagnosis)	No	228	35.09***
	Yes	34	58.82
Global Assessment of Functioning (GAF)	Median or below	131	45.80*
	Above median	131	30.53
Brief Symptom Inventory (BSI)	Median or below	135	33.33
	Above median	127	43.31
Paranoid symptoms (BSI sub-scale)	Median or below	137	29.93**
	Above median	125	47.20
Threat/control-override (TCO) symptoms	Median or below	134	31.34*
	Above median	128	45.31
Insight and Treatment Attitudes Questionnaire (ITAQ)	Median or below	135	42.96
	Above median	127	33.07
Psychiatric hospitalisations in the preceding year	Less than two	163	33.13*
	Two or more	99	46.46
Substance use	No use	112	26.79***
	Occasional use	60	36.67
	Problems	90	53.33
Medication-adherent in preceding four months	No	190	37.89
	Yes	72	38.89

Statistical significance (Fisher's exact test, two-tailed): † $P < 0.10$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Medication adherence

Medication adherence was measured from three interview sources: subject, case manager and collateral informant. Respondents were defined as non-adherent if they were prescribed psychotropic medications but reportedly never took them or only sometimes took them as prescribed (Swartz *et al*, 1998b).

Out-patient services utilisation

Out-patient services utilisation was obtained from service records in the information systems of participating community mental health centres. All service encounters for case management, medication, psychotherapy and other out-patient services were summed in a single index. Regular treatment was defined as three or more out-patient service encounters per month in the community (the median amount). This rate was adjusted for time spent hospitalised. For those receiving regular treatment according to this definition, the median number of out-patient service events per month was seven. For those not receiving regular treatment, the median number of events was one per month.

Analysis

Logistic regression analysis was used to examine the relative impact of sustained OPC and out-patient services utilisation on the incidence of violent behaviour, controlling for baseline violence history and relevant covariates. Odds ratios produced by this technique estimate the average change in the odds of a predicted event (any violence) associated with the presence of a risk factor. For independent variables measured on a continuous scale or ranking, the odds ratio indicates the change in event likelihood per unit change in the predictor.

Sample description

Demographic characteristics

As noted in Table 1, sample members were predominantly young to middle-aged adults of lower educational status and mostly not married and not cohabiting. The racial distribution of the sample was 66% African American, 33% non-Hispanic White and 1% other. Although the majority were city residents, a substantial proportion of the sample lived in rural areas and small towns. This sample was representative of the

population of patients admitted to state mental hospitals in North Carolina.

Clinical characteristics, substance misuse, and history of violence at baseline

The majority of the sample (68%) had diagnoses of psychotic disorders, whereas 28% had bipolar disorder and 4% had recurrent major depression. Thirteen per cent had a co-occurring diagnosis of a personality disorder. The majority of the sample had moderate functional impairment (GAF median score=47). Alcohol and drug use (57%), medication non-compliance (73%) and violent behaviour (51%) were common in the four months prior to hospitalisation. More than one-third (39%) had experienced two or more psychiatric hospital admissions during the preceding year. For a more extensive presentation of the sample and the characteristics of violent events preceding hospitalisation, see Swanson *et al* (1999).

RESULTS

Predictors of violence

Table 1 presents bivariate associations of baseline risk factors with any violent behaviour during the one-year follow-up. The risk of violence was significantly higher among subjects who were: young (under age 40 years); single; with low social support; residents of urban areas; those recently homeless; with greater functional impairment (GAF score <47); substance misusers; those with paranoid symptoms; TCO symptoms; and those with more than two hospital admissions in the prior year.

Key correlates of violent behaviour during the follow-up year included ongoing substance misuse problems and non-adherence with prescribed psychotropic medications. Fifty per cent of substance misusers were violent *v.* 26.5% of those who did not misuse substances (Fisher's exact test, two-tailed, $P < 0.001$); and 41.6% of those who did not adhere to prescribed medication regimens were violent *v.* 25.5% of those who took medications as prescribed (Fisher's exact test, two-tailed, $P < 0.05$).

Out-patient commitment and violence

Considering this study strictly as a randomised controlled trial, subjects with a baseline history of serious violence must be

excluded from analysis and all experimental subjects must be considered as a single group (without regard to the length of exposure to OPC). On this basis alone, the study found no significant difference in the prospective rate of violence between the two randomly assigned groups: 32.3% in the OPC group *v.* 36.8% in the control group (Fisher's exact test, one-tailed: $P = 0.292$; two-tailed: $P = 0.567$).

However, a problem with this test is that it excludes a key risk group to which OPC policy may be targeted specifically (the seriously violent group), and constricts the dependent variable by eliminating a substantial number of violent events from the analysis. Moreover, the test as specified cannot take into account varying amounts of exposure to the intervention (i.e. length of time actually spent under court-ordered treatment). Time on OPC varied considerably; it was not subject to experimental control, but depended on clinician's discretion in applying the legal criteria for renewal of expiring OPC orders. As mentioned earlier, this may have selected into the long-term OPC group a greater proportion of clients with a history of non-adherence to medication; this was indirectly associated with a higher baseline risk of violence. Clearly, however, insofar as this created a bias, its effect was conservative, that is, it would work *against* finding a positive effect for extended OPC in reducing violence risk.

Alternatively, when subjects with a history of serious violence were included in the analysis and the OPC intervention was re-defined as *receiving at least six months of court-ordered treatment*, the results were striking. The extended OPC group had a significantly lower incidence of violence during the year: 26.7% *v.* 41.6% (Fisher's exact test, one-tailed: $P = 0.049$; two-tailed: $P = 0.025$).

The six-month cut-off point for categorising extended OPC is somewhat arbitrary. However, there were good reasons for dichotomising this variable. The distribution of total OPC days was highly skewed, with all members of the control group receiving zero days of OPC. The majority of the randomised experimental and violent group subjects received less than six months of OPC, whereas a large cluster remained on OPC for the entire year of the study, giving the variable a bimodal distribution. Specifically, for those with less than six months of court-ordered treatment the mean number of days on

OPC was 76, whereas for those receiving extended OPC the mean number of days was 330. Using a dichotomous version of the intervention variable, a significant association was found between extended OPC and incidence of violence; this effect remained significant in multivariate analysis, as will be shown below.

We also re-analysed the data for high and low exposure to OPC separately for the randomised subjects and the non-randomised violent group. In the randomised study groups, control subjects did not differ significantly in their risk of violence from those with brief, discontinued OPC (36.8% *v.* 39.7%). However, the incidence of violence was significantly less (22.7%) among randomised OPC subjects who received *extended* court-ordered treatment (Fisher's exact test, one-tailed: $P = 0.043$; two-tailed: $P = 0.076$). Among the subjects who were violent at baseline, the effect of extended OPC was also seen at a level approaching statistical significance: among violent subjects with brief, discontinued OPC, 63.3% repeated their violent behaviour during the year of the study; of those receiving extended OPC, only 37.5% were violent (Fisher's exact test, one-tailed: $P = 0.086$; two-tailed: $P = 0.126$).

Multivariate analysis

Staged logistic regression analysis with stepwise inclusion was conducted to test the effect of OPC on reduced violence in the context of relevant covariates. A control variable was included to hold constant the effect of baseline history of violence and initial assignment to the violent OPC group. An initial model (see Table 2, Model 1) showed that extended OPC was associated with significantly lower odds of any violent behaviour during the year of the study, controlling for baseline history of violence (odds ratio=0.42; $P < 0.05$). In Model 2, being young, single, and having a low degree of perceived social support were identified as significant demographic risk factors. In Model 3, among the clinical variables substance misuse was a significant predictor of violence. Controlling for the significant demographic variables and substance misuse, respondents who received more than 180 days of OPC were only about one-third as likely to commit a violent act during the year as their counterparts in the control group (odds ratio=0.35; $P < 0.01$). Those receiving 1–179 days of OPC did not differ

from the control group with respect to risk of violence.

A cross-sectional analysis of baseline data from this same study showed a significant association between history of violence and history of medication non-adherence combined with substance misuse (Swartz *et al*, 1998b). This finding was replicated with the longitudinal follow-up data controlling for extended OPC and significant covariates. Risk of violence was found to be three times greater among subjects who misused substances and did not take medications as prescribed during follow-up (odds ratio=3.19; $P < 0.01$); however, risk was not significantly elevated among respondents with only one of these problems alone.

Based on this finding of an interaction effect of substance misuse and non-compliance, two composite variables were coded for subsequent analysis: clinical risk status at baseline, and change in clinical risk status at follow-up. Subjects whose status on both substance misuse and medication adherence improved from baseline received the highest change scores, whereas those who deteriorated or maintained a poor status received lower scores accordingly. Change scores ranged from -3 to +4 (mean=1.2, s.d.=1.7). In bivariate analysis, subjects who received extended OPC achieved significantly better mean scores on the index (1.8 *v.* 1.1; mean square=14.2; $F=4.79$; $P < 0.05$). Subjects who remained free of violent behaviour

also were found to have significantly better scores (1.53 *v.* 0.76; mean square=36.7; $F=12.76$; $P < 0.01$).

The staged multivariate analysis shown in Table 3 examines whether OPC interacts with the provision of out-patient services to reduce the risk of violent behaviour and, if so, whether OPC prevents violence indirectly by means of reducing substance misuse and improving medication adherence. Model 1 shows that extended OPC *alone* did not significantly reduce risk. Similarly, receiving frequent services *alone* was not associated with less violence. Rather, the combination of both variables – at least 180 days of OPC with an average of three or more out-patient visits per month in the

Table 2 Predictors of violent behaviour during the year after hospital discharge: effects of sustained out-patient commitment (OPC) controlling for baseline risk factors

Domain	Independent variable	Model 1	Model 2	Model 3
		Odds ratio (95% CI)	Odds ratio (95% CI)	Odds ratio (95% CI)
Prior history of violence Intervention	None/fights/serious violence	1.954*** (1.341–2.846)	2.083*** (1.373–3.161)	1.915** (1.262–2.906)
	Zero days OPC (control)	1.000	1.000	1.000
	1–179 days OPC	1.049 NS (0.563–1.954)	0.995 NS (0.499–1.984)	0.986 NS (0.500–1.945)
	180 days or more OPC	0.416* (0.197–0.880)	0.417* (0.185–0.940)	0.347* (0.152–0.792)
Demographic characteristics	Younger age (under 40 years)		2.440** (1.375–4.332)	2.129* (1.196–3.791)
	Single <i>v.</i> married/cohabiting		2.298* (1.054–5.012)	2.517* (1.165–5.439)
	Social support		0.894*** (0.842–0.950)	0.895*** (0.843–0.951)
	Urban <i>v.</i> rural resident		1.695† (0.927–3.102)	
	African American		1.708† (0.927–3.147)	
	Male <i>v.</i> female		NS	
	Income		NS	
	Education		NS	
	Homeless		NS	
	Victim of crime		NS	
Clinical characteristics	Substance misuse problems			2.360** (1.291–4.314)
	Diagnosis: psychotic <i>v.</i> affective			NS
	Personality disorder (Axis 2 diagnosis)			NS
	Functional impairment (GAF)			NS
	Brief Symptom Inventory (BSI)			NS
	Paranoid symptoms (BSI sub-scale)			NS
	Threat/control override symptoms			NS
	Insight into illness (ITAQ)			NS
	Prior psychiatric hospitalisations			NS
	Medication adherence			NS
Model statistics	Number of observations	262	245	256
	Degrees of freedom	2	6	6
	-2 log likelihood χ^2	19.610***	57.448***	56.311***
	Rank correlation predicted/observed	0.650	0.770	0.761

Variables in each domain were entered stepwise and retained in the subsequent stage model if $P < 0.10$. Statistical significance: NS=not selected/not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

community – showed a significant effect in reducing violence (odds ratio=0.37; $P<0.05$). In this model, the predicted probability of any violent behaviour was cut in half from 48% to 24%, attributable to extended OPC and regular out-patient services provision.

Model 2 in Table 3 shows that combined improvement in medication adherence and substance use/misuse status was a significant predictor of reduced violent behaviour: about 20% reduction with each unit improvement in clinical risk (odds ratio=0.81; $P<0.05$). This model provides evidence of an indirect effect of the intervention, as mentioned above. With the clinical risk change score in the equation, the direct effect of OPC and services was suppressed from an odds ratio of 0.36 ($P<0.05$) in Model 1 to an odds ratio of 0.41 ($P<0.10$) in Model 2, and the fit of the model improved significantly.

Finally, Model 3 specifies a reduced set of predictors using stepwise selection of variables. Based on this model, subjects who received the high intervention (sustained OPC and regular services), who concurrently remained free of substance misuse and who took medications as prescribed during the year of the study had the lowest likelihood of any violence (13% predicted probability). In contrast, those who did not receive the high intervention, who continued to misuse substances and who stopped taking their

medications had a 53% predicted probability of violent behaviour during the year.

DISCUSSION

This study examined jointly the effectiveness of OPC and the impact of key risk factors associated with violent behaviour among persons with psychotic or major mood disorders. In a multivariate model that tested a range of socio-demographic and clinical characteristics, a small set of baseline risk factors – being young (under age 40 years), single, having low social support, and misusing substances – were identified prospectively as significant predictors of violent behaviour in the ensuing year. Consistent with prior analysis (Swartz *et al*, 1998b), the study also found that the combination of co-occurring substance misuse and non-adherence with medications was strongly associated with violence.

Sustained OPC combined with regular community-based service

The key finding is that extended OPC (more than six months) combined with regular out-patient services utilisation (three or more events per month) produced a significant decrease in the incidence of violence. Neither the court order nor services *alone* was effective in reducing violence. However, with regular service use, the presence

of extended OPC reduced the probability of violence from 48% to 24%.

Improved medication adherence and diminished substance misuse

The study provides evidence for an important mechanism by which OPC may exert its effect in preventing violent behaviour. Respondents who received extended OPC with regular out-patient services achieved, on average, the highest positive (improved) change scores on medication adherence and substance use. In turn, those with high change scores in both of these risk factors combined had a significantly reduced incidence of violent behaviour during the year of follow-up. A multivariate model suggests that the benefit of OPC as described is at least partly an indirect effect attributable to change in these intervening clinical risk factors.

Deviation from randomised design

The study design deviates in two ways from a strict randomised controlled trial. First, the sample included a subgroup of subjects with a recent history of serious violent behaviour who could not be assigned randomly to the initial control group (outright release from OPC). To exclude these subjects altogether would have meant that the study findings could not be generalised to a crucially important sub-population to

Table 3 Predictors of behaviour during the year after hospital discharge: effects of sustained out-patient commitment (OPC) and regular services controlling for baseline characteristics and improvement in combined clinical risk factors (substance misuse and medication adherence)

Domain	Independent variable	Model 1	Model 2	Model 3
		Odds ratio (95% CI)	Odds ratio (95% CI)	Odds ratio (95% CI)
Prior history of violence	None/fights/serious violence	1.976*** (1.336–2.923)	1.99*** (1.334–2.967)	1.912*** (1.292–2.831)
Intervention	No extended OPC/no regular services (comparison)	1.000	1.000	
	Regular services without extended OPC	0.613 (0.205–1.831)	0.655 (0.213–2.015)	
	Extended OPC without regular services	1.570 (0.824–2.990)	1.508 (0.786–2.895)	
	Extended OPC with regular services	0.365* (0.139–0.958)	0.411† (0.154–1.095)	0.358* (0.145–0.883)
Baseline demographics	Younger age (under 40 years)	2.112* (1.182–3.774)	1.909* (1.056–3.451)	1.933* (1.074–3.480)
	Single v. married/cohabiting	2.400* (1.103–5.223)	2.381* (1.078–5.258)	2.454* (1.119–5.381)
	Social support	0.907** (0.853–0.963)	0.905** (0.852–0.962)	0.899*** (0.847–0.955)
Baseline clinical risk score	Substance misuse (SM)+medication non-adherence	1.390* (1.045–1.849)	1.385* (1.036–1.852)	1.376* (1.032–1.835)
Change in clinical risk score	Improvement in SM status+medication adherence		0.811* (0.682–0.965)	0.801* (0.675–0.951)
Model statistics	Number of observations	258	258	258
	Degrees of freedom	8	9	7
	–2 log likelihood χ^2	58.529***	64.269***	61.325***
	Improvement in fit: Model 2 v. Model 1 ($\chi^2/1$ d.f.)		5.800*	
	Rank correlation predicted/observed	0.762	0.778	0.772

Statistical significance: † $P<0.10$; * $P<0.05$; ** $P<0.01$; *** $P<0.001$.

which OPC policy may be targeted. However, random assignment was not feasible for these subjects; the Institutional Review Board, the District Court judges and the hospital psychiatrists overseeing these patients' care and research participation would not authorise rescinding their OPC orders at baseline – citing practice liability and other concerns. Nevertheless, over half of these subjects *did not remain* on OPC, and thus were unexposed to court-ordered treatment during a large proportion of the period of observation. Hence, the non-renewal of many of these orders allowed an informative comparison of rates of violent recidivism between those remaining under OPC for more than six months (average 330 days) and those receiving less than six months of OPC (average 76 days).

This leads to a second necessary deviation from a strict randomised controlled design. Amount of time on OPC was not random or controlled experimentally, but varied as clinicians applied the legal criteria for renewal of OPC orders. Potentially, this could have led to a biased conclusion (i.e. attributing a positive intervention effect to subjects who might have been less violent anyway because of pre-existing lower risk). However, this could only be a problem if higher-risk subjects were less likely to get renewal of their court order. In fact, the legal criteria for OPC work in the opposite direction.

Possible conservative bias favouring not finding an effect of OPC on reducing violence

Renewal of the court order required a second determination (by a psychiatrist and the court) that the respondent would predictably become dangerous (or 'gravely disabled') without treatment and predictably would not comply with treatment. At the end of the initial OPC period (up to 90 days), each case was re-evaluated systematically. Prompting notices were sent to clinicians, reminding them that a subject's OPC order was about to expire and summarising the OPC criteria for easy reference. If at that point the psychiatrist and the court concluded that the respondent was no longer likely to become dangerous without treatment or, even if so, would comply *voluntarily* with treatment, then the legal criteria for OPC were not satisfied and the order could not be renewed. (Beyond the legal criteria, common sense suggests that clinicians would not cull out clients

CLINICAL IMPLICATIONS

- An extended period of involuntary out-patient commitment (six months or more) may significantly reduce the risk of violent behaviour in persons with severe mental illness.
- Involuntary out-patient commitment must be supported by frequent mental health service visits (three or more monthly visits on average) in order to have a significant effect in managing violence risk.
- Out-patient commitment reduces violence risk at least in part by improving medication adherence while diminishing substance misuse; hence, treatment planning for out-patient-committed persons with a history of violence should focus on these intervening clinical issues.

LIMITATIONS

- Length of time on out-patient commitment was a key variable in the intervention, but could not be randomly assigned. Legal criteria for renewal of out-patient commitment would seem to prevent selection of lower-risk subjects for longer exposure to court-ordered treatment. Still, unknown selection factors may have affected duration of out-patient commitment.
- Persons with a documented history of serious violent behaviour upon entry into the study could not be assigned to control group (released from out-patient commitment outright); hence, analysis of the effectiveness of out-patient commitment in this group is limited to comparisons between those with shorter *v.* longer periods of court-ordered treatment in the community.
- Out-patient service intensity was not controlled by the study, but varied according to clinical need and unknown factors. Perceived risk may have stimulated service provision, just as treatment may have prevented violence; causation is difficult to establish.

JEFFREY W. SWANSON, PhD, MARVIN S. SWARTZ, MD, H. RYAN WAGNER, PhD, BARBARA J. BURNS PhD, Services Effectiveness Research Program, Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC; RANDY BORUM, PsyD, Department of Mental Health Law and Policy, Florida Mental Health Institute, University of South Florida, Tampa, FL; VIRGINIA A. HILDAY, PhD, Department of Sociology, North Carolina State University, Raleigh, NC, USA

Correspondence: Jeffrey Swanson, Assistant Professor in Psychiatry and Behavioral Sciences, Duke University Medical Center, Box 3071, Brightleaf Square Suite 23-A, Durham, NC 27710, USA. Tel: (919) 682 4827; Fax: (919) 682 1907; e-mail: Jeffrey.Swanson@Duke.edu

(First received 19 May 1999, final revision 8 December 1999, accepted 9 December 1999)

assessed to be at highest risk of violence and selectively *not* renew their OPC orders.)

Anecdotally, when clinicians were asked by research staff members to state their reasons for not renewing an OPC order, the most typical answer was that the patient had been compliant with treatment and was doing reasonably well at that time; hence, continuing a court order was

not seen as legally justifiable. Empirically, respondents who had been mostly compliant with medications in the four months prior to hospitalisation were significantly less likely to receive extended OPC after their initial court order expired; and medication compliance was indirectly associated with lower risk of violence in these data (Swartz *et al*, 1998b). In summary, if bias existed in the selection of subjects for

longer periods of OPC, then such a bias would seem to work *against* finding that extended OPC lowers the risk of violence. Thus, our results may understate the true impact of OPC in preventing violent behaviour.

Responsibility of mental health service systems using OPC

These results show that OPC may be an effective strategy for reducing community violence among people with SMI. However, if service systems intend to use OPC to this end, they must be prepared to monitor clients for sustained periods. Most importantly, they must make available adequate services to address the complex needs of persons with SMI and (often) substance misuse comorbidity, who lack adequate social resources and experience persistent difficulties in complying with out-patient treatment on their own. With such provisos, OPC may assist mental health agencies in reducing violent behaviour and costly institutional recidivism, and may significantly improve the lives of persons suffering from severe and persistent mental illness.

ACKNOWLEDGEMENTS

The research presented here was supported by two grants from the National Institute of Mental Health: Effectiveness of Involuntary Outpatient Commitment (R01-MH48103-05); and Program on Services Research for People with Severe Mental Disorders (P50-MH51410-02). The authors gratefully acknowledge the support and cooperation of many state mental health personnel, judges, clinicians, administrative staff and law officers from Durham, Vance, Franklin, Granville, Warren, Orange, Person, Chatham and Guilford counties in North Carolina and Linda George, PhD, who helped design and participated in this study.

REFERENCES

- Angermeyer, M. & Matschinger, H. (1996)** The effect of violent attacks by schizophrenic persons on the attitude of the public towards the mentally ill. *Social Science and Medicine*, **43**, 1721–1728.
- Borum, R., Swartz, M. & Swanson, J. (1996)** Assessing and managing violence risk in clinical practice. *Journal of Practical Psychiatry and Behavioral Health*, **2**, 205–215.
- Cuffel, B. (1997)** Disruptive behaviour and the determinants of costs in the public mental health system. *Psychiatric Services*, **48**, 1562–1566.
- Derogatis, L. & Melisaratos, N. (1983)** The brief symptom inventory: a brief report. *Psychological Medicine*, **13**, 595–605.
- Endicott, J., Spitzer, R., Fleiss, J., et al (1976)** The global assessment scale: a procedure for measuring overall severity of psychiatric disturbance. *Archives of General Psychiatry*, **33**, 766–771.
- Fernandez, G. & Nygard, S. (1990)** Impact of involuntary outpatient commitment on the revolving-door syndrome in North Carolina. *Hospital and Community Psychiatry*, **41**, 1001–1004.
- Geller, J., Grudzinskas, A., McDermeit, M., et al (1998)** The efficacy of involuntary outpatient treatment in Massachusetts. *Administration and Policy in Mental Health*, **25**, 271–285.
- George, L. K., Blazer, D. G., Hughes, D. C., et al (1989)** Social support and the outcome of major depression. *British Journal of Psychiatry*, **154**, 478–485.
- Link, B., Cullen, F., Frank, J., et al (1987)** The social rejection of former mental patients: understanding why labels matter. *American Journal of Sociology*, **92**, 1461–1500.
- , Stueve, A. & Phelan, J. (1998)** Psychotic symptoms and violent behaviors: probing the components of 'threat/control-override' symptoms. *Social Psychiatry and Psychiatric Epidemiology*, **33**, 55–60.
- McEvoy, J., Apperson, L., Applebaum, P., et al (1989)** Insight into schizophrenia: its relationship to acute psychopathology. *Journal of Nervous and Mental Disease*, **177**, 43–47.
- Monahan, J. (1992)** Mental disorder and violent behavior: perceptions and evidence. *American Psychologist*, **47**, 511–521.
- & Steadman, H. (1994)** *Violence and Mental Disorder: Developments in Risk Assessment*. Chicago, IL: University of Chicago Press.
- Mullen, P. (1997)** A reassessment of the link between mental disorder and violent behaviour, and its implications for clinical practice. *Australian and New Zealand Journal of Psychiatry*, **31**, 3–11.
- Mulvey, E., Geller, J. & Roth, L. (1987)** The promise and peril of involuntary outpatient commitment. *American Psychologist*, **42**, 571–584.
- Silver, E., Mulvey, E. P. & Monahan, J. (1999)** Assessing violence risk among discharged psychiatric patients: toward an ecological approach. *Law and Human Behavior*, **23**, 237–255.
- Simon, R. (1998)** Psychiatrists' duties in discharging sicker and potentially violent inpatients in the managed care era. *Psychiatric Services*, **49**, 62–67.
- Stefan, S. (1987)** Preventive commitment: the concept and its pitfalls. *Mental and Physical Disability Law Reporter*, **11**, 288–302.
- Swanson, J., Estroff, S., Swartz, M., et al (1997a)** Violence and severe mental disorder in clinical and community populations: the effects of psychotic symptoms, comorbidity, and lack of treatment. *Psychiatry*, **60**, 1–22.
- , Swartz, M., George, L., et al (1997b)** Interpreting the effectiveness of involuntary outpatient commitments: a conceptual model. *Journal of the American Academy of Psychiatry and Law*, **25**, 5–16.
- , Borum, R., Swartz, M., et al (1999)** Violent behaviour preceding hospitalization among persons with severe mental illness. *Law and Human Behavior*, **23**, 185–204.
- Swartz, M., Swanson, J., Hiday, V., et al (1998a)** Taking the wrong drugs: substance abuse, medication nonadherence, and violence in severely mentally ill individuals. *Social Psychiatry and Psychiatric Epidemiology*, **33**, 75–80.
- , —, —, et al (1998b)** Violence and severe mental illness: the effects of substance abuse and nonadherence to medication. *American Journal of Psychiatry*, **155**, 226–231.
- , —, Wagner, H., et al (1999)** Can involuntary outpatient commitment reduce hospital recidivism? Findings from a randomised trial in severely mentally ill individuals. *American Journal of Psychiatry*, **156**, 1968–1975.
- Torrey, E. & Kaplan, R. (1995)** A national survey of the use of outpatient commitment. *Psychiatric Services*, **46**, 778–784.